



THE ANNALS OF THORACIC SURGERY



Post-Traumatic Complex Aortic Arch Pseudoaneurysm With Aorto-Innominate Vein Fistula

Michael Malyshev, Alexander Safuanov, Dmitry Siniukov, Dmitry Borovikov, Anton
Malyshev and Andrew Rostovykh

Ann Thorac Surg 2009;88:1025

DOI: 10.1016/j.athoracsur.2008.12.092

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://ats.ctsnetjournals.org/cgi/content/full/88/3/1025>

The Annals of Thoracic Surgery is the official journal of The Society of Thoracic Surgeons and the Southern Thoracic Surgical Association. Copyright © 2009 by The Society of Thoracic Surgeons. Print ISSN: 0003-4975; eISSN: 1552-6259.

Post-Traumatic Complex Aortic Arch Pseudoaneurysm With Aorto-Innominate Vein Fistula

Michael Malyshev, MD, PhD, Alexander Safuanov, MD, Dmitry Siniukov, MD, Dmitry Borovikov, MD, Anton Malyshev, MD, and Andrew Rostovkykh, MD

Center of Cardiac Surgery, Chelyabinsk, Russia




Fig 1.




Fig 2.




Fig 3.

A 43-year-old man was admitted to our institute in cardiac failure 12 years after he had sustained a stab wound in the suprasternal notch. A thoracic aortogram demonstrated a communication between the distal aortic arch and the left innominate vein (LIV) within consecutive double pseudoaneurysms (Figs 1 and 2; white and black arrows respectively). Figure 3 shows the venous phase image of the aortogram that showed the arteriovenous (AV) fistula and enlarged left innominate vein.

The patient was operated on through a median sternotomy with hypothermic (20°C) circulatory arrest. The distal aortic arch and the calcified pseudoaneurysms were dissected free as much as possible before initiation of cardiopulmonary bypass. The direct digital pressure on the venous end of communication was used to prevent systemic hypoperfusion during the patient's cooling. Once the temperature of 20°C was reached, cardiopulmonary bypass was stopped, and the lumen of the pseudoaneurysm communicated with left innominate vein was opened (Fig 4), and the fragmentary was removed. The pseudoaneurysm adjacent to the aortic arch was also removed with the aortic wall. The defects of the aortic arch and of the left innominate vein were repaired using a xenopericardial patch and a direct suture, correspondingly. After the air was removed from the aorta, and cardiopulmonary bypass was recommenced, the patient was re-warmed. He was still asymptomatic at his postoperative 8-month follow-up.

Address correspondence to Dr Malyshev, Center of Cardiac Surgery, PO Box 5977, Chelyabinsk, 454047, Russian Federation; e-mail: cardiosur@chel.surnet.ru.




Fig 4.

The identified lesion is a rare case, and its anatomical forms are variable. The safe universal surgical repair has been developed after the first introduction by Borst and colleagues [1] in 1964. The optimal surgical technique includes cardiopulmonary bypass with hypothermic circulatory arrest through the median sternotomy, use or preparation of the femoral vessels for cannulation, prevention of the inadequate systemic perfusion by digital compression of the fistula or clamping of the left innominate vein [2].

References

1. Borst HG, Schaudig A, Rudolph W. Arteriovenous fistula of the aortic arch: repair during deep hypothermia and circulatory arrest. *J Thorac Cardiovasc Surg* 1964;48:443.
2. Reddi A, Madansein R, Mathivha T. Traumatic arteriovenous fistula of ascending aorta and left innominate vein. *Ann Thorac Surg* 2003;75:1637-40.

Post-Traumatic Complex Aortic Arch Pseudoaneurysm With Aorto-Innominate Vein Fistula

Michael Malyshev, Alexander Safuanov, Dmitry Siniukov, Dmitry Borovikov, Anton Malyshev and Andrew Rostovykh

Ann Thorac Surg 2009;88:1025

DOI: 10.1016/j.athoracsur.2008.12.092

Updated Information & Services

including high-resolution figures, can be found at:
<http://ats.ctsnetjournals.org/cgi/content/full/88/3/1025>

References

This article cites 2 articles, 1 of which you can access for free at:
<http://ats.ctsnetjournals.org/cgi/content/full/88/3/1025#BIBL>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Great vessels

http://ats.ctsnetjournals.org/cgi/collection/great_vessels

Permissions & Licensing

Requests about reproducing this article in parts (figures, tables) or in its entirety should be submitted to:

<http://www.us.elsevierhealth.com/Licensing/permissions.jsp> or
email: healthpermissions@elsevier.com.

Reprints

For information about ordering reprints, please email:
reprints@elsevier.com



THE ANNALS OF THORACIC SURGERY

